

MaxiFlora

Commercial Fertilizer

0-0-6



Guaranteed analysis:

Soluble Potash (K₂O)6.00%
Sulfur (S)2.00%

Derivation:

Potash and Sulfur derived from Potassium Sulfate.

Weight per gallon = 8.80 Lbs.
Kilograms per liter = 1.05 Kg.

Handling:

Shake or stir before use. Follow appropriate safety procedures. In case of accidental exposure, flush with plenty of water. Product is non-hazardous. Storage for more than one year is not recommended.

Incompatibilities:

Always jar test newly attempted mixes. No known incompatibilities.

Directions for use:

MaxiFlora is a supplemental nutrient for use with foliar spray applications. Properly timed applications may aid in improved crop nutrition. Use MaxiFlora with a soil fertility program and good cultural practices. The use of soil and tissue testing is recommended to determine the need for additional nutrients. Use 0.5 - 2.0 gallons per acre during the growing season, as needed.

Directions for application:

Application instructions: SHAKE WELL BEFORE USE. Apply with enough water to ensure thorough coverage.

For aircraft, use 5-10 gallons of water per acre and for ground applications use 10-250 gallons per acre. For best results apply MaxiFlora in the early mornings or late evening with a quality surfactant.

General use guidelines:

MaxiFlora may be applied to all flowering crops at 1-4 quarts per application and up to 4 applications per season. This includes, but is NOT limited to the following crops:

Crop	Stage	Rate	Application*
Apples, Pears	Bud Break	1 qt.	Do not apply to mature fruit
	First Green Leaf	1.5 qts.	
	3 week interval	1 qt.	
Apricot, Cherry, Peach, Plum, Nectarine	Bud Break	1.5 qts.	Do not apply to mature fruit
	First green leaf	2 qts.	
	3 week intervals	1.5 qts.	
Beans	Fourth true leaf	2 qts.	Make 4 appl.
	2 week intervals	1-2 qts.	
Broccoli, Cauliflower	Emergence	2 qts.	Directed spray Make 4 appl.
	2-3 week intervals	2 qts.	
Carrots	4-6 " Height 15-20 day interval	2 qts. 1-2 qts.	Make 4 appl.
Citrus	Pre bloom 2 week interval	2 qts. 2 qts.	Do not apply to mature fruit
Corn	6-16 " Height 2 week intervals	2 qts. 1.5 qts.	
Cotton	8-12 " Height	1-2 qts.	May be applied with pesticides
	2-3 week interval	1-2 qts.	
	Pre-cutout	2 qts.	
Cucumbers, Squash	3-4 leaf	1.1.5 qts.	Continue until 3 before harvest
	7-10 day interval	1 qt.	

Garlic	Leek stage	1.5 qts.	
	20-30 days later	1.5 qts.	
	15-20 days later	1.5 qts.	
Grapes	First leaf	1.5 qts.	
	Bud initiation	1 qt.	
	2-3 week interval	1-2 qts.	
Leafy vegetables Lettuce, Spinach	3-4 leaf	1-2 qts.	Apply after cutting
	2-3 week interval	1-2 qts.	
Melons	3 leaf	1-2 qts.	Continue until 3 weeks before harv.
	2-3 week interval	1-2 qts.	
Onions	3 leaf	2 qts.	
	2 week interval	2 qts.	
Potato, Radish	6-8 " height	1.5 qts.	
	2-3 week intervals	1.5 qts.	
Strawberry	First Bloom 2-3 week interval	1.5 qts. 1.5 qts.	Continue through harvest season.
Sugarbeet	2-3 week interval	1-2 qts.	Make 4 appl.
Tomatoes	8-10 " Height	2 qts.	Make 4 appl.
	2 week interval	2 qts.	
Wheat, Barley	End of tillering beginning of stem elongation	2 qts.	

* Always jar test newly attempted mixes.

Notice to user:

Call 1-800-USA-BIO1 for technical assistance concerning use rates for crops not mentioned in this label.

Manufactured by,
BioScientific, Inc.

Avondale, Arizona